arXiv:1410.4059v1 [math.GR] 15 Oct 2014

GENERATION OF REAL MUTUALLY UNBIASED BASES AS POWERS OF A REAL ORTHOGONAL MATRIX OF ODD FINITE ORDER

ROD GOW

Abstract. Let $q = 2^r$, where r is a positive integer. We show by group representation theory and hermitian geometry over a finite field that there exists a $q^2 \times q^2$ real orthogonal matrix D, say, of multiplicative order $q+1$ whose powers generate $q+1$ real pairwise mutually unbiased bases in $\mathbb{R}^{q^2}$. This means that the scaled powers $qD, \ldots, qD^q$ are q different real Hadamard matrices. We also show that each diagonal entry of these q Hadamard matrices equals -1.

1. Introduction

Let d be a positive integer and let $\mathbb{R}^d$ denote the Hilbert space of dimension d over the field $\mathbb{R}$ of real numbers. Let $g : \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}$ denote a positive definite symmetric bilinear form. Let $B_1 = \{ u_1, \ldots, u_d \}$ and $B_2 = \{ v_1, \ldots, v_d \}$ be bases of $\mathbb{R}^d$ that are orthonormal with respect to g. We say that the bases B_1 and B_2 are *mutually unbiased* if

$$g(u_i, v_j)^2 = \frac{1}{d}$$

for $1 \leq i, j \leq n$. We may identify the orthonormal bases B_1 and B_2 with $d \times d$ real orthogonal matrices O_1 and O_2, say. Then the condition that the bases are mutually unbiased is equivalent to saying that each entry of $O_1 O_2^{-1}$ equals $\pm 1/\sqrt{d}$.

It is known that the maximum number of mutually unbiased bases in $\mathbb{R}^d$ is $1 + \frac{d}{2}$, but a combination of theory and experimental evidence suggests that this upper bound is rarely met. We note, however, that the upper bound is met when $d = 2^{2r}$, where r is a positive integer, [3]. The introduction to the paper [2] provides useful information on the problem of determining the maximum number of real mutually unbiased bases.

When investigating several mutually unbiased bases, we may assume that the basis B_1 is the standard basis of $\mathbb{R}^d$, corresponding to the identity matrix I_d. Then if the basis B_i corresponds to the orthogonal matrix O_i, B_i is mutually unbiased with respect to B_1 if and only if O_i is an Hadamard matrix scaled by $1/\sqrt{d}$. Similarly, B_i and B_j are mutually unbiased if and only if $O_i O_j^{-1}$ is also an Hadamard matrix scaled by $1/\sqrt{d}$. This fact explains why if $d \neq 2$, and 4 does not divide d, there do not exist two mutually unbiased bases in $\mathbb{R}^d$.

The purpose of this paper is to show that when $d = 2^{2r}$, we can find a $d \times d$ real orthogonal matrix D, say, of multiplicative order $2^r + 1$, whose $2^r + 1$ powers D, $D^2, \ldots, D^{2^r+1} = I_d$, define $2^r + 1$ mutually unbiased bases in $\mathbb{R}^d$. Thus the scaled

School of Mathematical Sciences, University College, Belfield, Dublin 4, Ireland, *E-mail address:* rod.gow@ucd.ie.

matrices 2^rD, ..., $2^rD^{2^r}$ are 2^r different Hadamard matrices. We also show that the diagonal entries of these Hadamard matrices all equal -1.

Our proof is group-theoretic, and makes use of elementary hermitian geometry over a finite field, as well as basic ideas in complex and real representation theory. A consequence of this approach is that we do not have an explicit description of D and its powers, beyond the diagonal property mentioned above. Our proof is in the spirit of our previous demonstration that when $d = 2^r$, there exists a $d \times d$ complex unitary matrix of multiplicative order $d+1$ whose $d+1$ different powers constitute $d+1$ mutually unbiased bases in the complex Hilbert space $\mathbb{C}^d$, [4]. We recall that $d+1$ is the maximum number of mutually unbiased bases in $\mathbb{C}^d$.

We have not been able to solve the problem of deciding if there exists a $2^{2r} \times 2^{2r}$ real orthogonal matrix of multiplicative order $2^{2r-1}+1$ whose different powers provide us with $2^{2r-1}+1$ mutually unbiased bases in $\mathbb{R}^{2^{2r}}$, the maximum number possible. If such a matrix exists, we would have the real analogue of the complex phenomenon just described. Of course, such a matrix exists when $r = 1$, and it is decribed in our paper of 2007.

2. Hermitian Geometry And The Unitary Group

The theory we require exists in various parts of the research literature but to try to keep this exposition reasonably elementary and self-contained, we will prove most of what we need to know.

Let $q = 2^r$, where r is a positive integer. Let $\mathbb{F}_{q^2}$ denote the finite field of order q^2. Let V be a vector space of dimension two over $\mathbb{F}_{q^2}$ and let $f : V \times V \to \mathbb{F}_{q^2}$ denote a non-degenerate hermitian form. Thus, we have

$$\begin{aligned} f(u+w,v) &= f(u,v)+f(w,v) \\ f(\lambda u,v) &= \lambda f(u,v) \\ f(v,u) &= f(u,v)^q \end{aligned}$$

for all u, v and w in V and all λ in $\mathbb{F}_{q^2}$. Furthermore, the non-degeneracy of f means that the only element u of V that satisfies $f(u,w) = 0$ for all $w \in V$ is the zero element.

Let G denote the set $V \times \mathbb{F}_{q^2}$. We define a multiplication on G by putting

$$(u,a)(v,b) = (u+v, f(u,v)+a+b)$$

for all ordered pairs (u,a) and (v,b) in G. It is straightforward to see that this multiplication is associative and makes G into a finite group of order q^6. The identity element is $(0,0)$ and the inverse of (u,a) is $(u, f(u,u)+a)$. The fact that f is non-degenerate implies that the center $Z(G)$ of G consists of all elements $(0,a)$, where $a \in \mathbb{F}_{q^2}$, and is thus elementary abelian of order q^2.

We set $x = (u,a)$, $y = (v,b)$ and let $[x,y]$ denote the commutator $x^{-1}y^{-1}xy$. We find that

$$[x,y] = (0, f(u,v)+f(v,u)).$$

Since f is non-degenerate and hermitian, the elements $f(u,v)+f(v,u)$ run through the subfield $\mathbb{F}_q$ of $\mathbb{F}_{q^2}$. Thus the commutator sugbroup G' of G, which is generated by the commutators $[x,y]$, consists of all elements of the form $(0,a)$, where a runs over $\mathbb{F}_q$. We deduce that $|G'| = q$.

We again set $x = (u,a)$ and assume now that $u \neq 0$. It is easy to check that the centralizer of x consists of all elements (v,b), where $v \in V$, b is an arbitary

element of $\mathbb{F}_{q^2}$, and $f(u,v) = f(v,u)$. Since $f(v,u) = f(u,v)^q$, if we consider V as a four-dimensional vector space over $\mathbb{F}_q$, we see that the set of all such v is a three-dimensional $\mathbb{F}_q$-subspace of V. Thus the centralizer of x has order q^5.

Next, we consider the irreducible complex characters of G.

Lemma 1. *The group G has exactly q^5 irreducible complex characters of degree 1. All other irreducible complex characters of G have degree q^2 and there are $q^2 - q$ of them. The irreducible characters of degree q^2 vanish outside $Z(G)$.*

Proof. Since the commutator subgroup G' has order q, G has exactly $|G : G'| = q^5$ different irreducible characters of degree 1. Now let x be any non-central element of G. The second orthogonality relation of group characters implies that, as the centralizer of x has order q^5,

$$q^5 = \sum |\chi(x)|^2,$$

where the sum extends over all irreducible characters χ of G. The characters of degree 1 each contribute 1 to this sum, and since there are q^5 of them, we deduce that $\psi(x) = 0$ for any irreducible character ψ of degree greater than 1.

Taking such an irreducible character ψ of G, Schur's Lemma implies that $|\psi(z)| = \psi(1)$ for all elements z of $Z(G)$ and hence we obtain by the first orthogonality relation of group characters that

$$|G| = q^6 = \sum |\psi(z)|^2 = |Z(G)|\psi(1)^2 = q^2\psi(1)^2,$$

where the sum extends over all elements z of $Z(G)$. We deduce that $\psi(1) = q^2$, as required.

Finally, we have

$$|G| = q^6 = \sum |\chi(1)|^2,$$

where the sum extends over all irreducible characters χ of G. Again, the characters of degree 1 each contribute 1 to this sum, and each irreducible character of degree q^2 contributes q^4 to the sum. Thus if there are r irreducible characters of degree q^2, we obtain

$$q^6 = q^5 + rq^4$$

It follows that $r = q^2 - q$, as claimed above. □

The next aspect of the representations of G that we wish to establish is that all of its complex irreducible characters are the characters of representations defined over the real field $\mathbb{R}$. To this end, given an irreducible complex character χ of G, we recall that its Frobenius-Schur indicator $\nu(\chi)$ is defined by the formula

$$\nu(\chi) = \frac{1}{|G|} \sum_{g \in G} \chi(g^2).$$

The general theory invented by Frobenius and Schur shows that $\nu(\chi)$ takes one of the three values 1, -1, and 0. We have $\nu(\chi) = 1$ if χ is a character of a representation of G defined over $\mathbb{R}$. We have $\nu(\chi) = -1$ if χ is real-valued but χ is not the character of a representation of G defined over $\mathbb{R}$. Finally, $\nu(\chi) = 0$ if χ is not real-valued. See, for example, [5], Chapter 4.

For our purposes, the following formula is useful. See [5], Corollary 4.6. We recall here that an *involution* in G is an element of order two.

Lemma 2. *Suppose that the number of involutions in G is t. Then we have*

$$\sum_{\chi} \nu(\chi)\chi(1) = t + 1,$$

where the sum is over all irreducible characters χ of G.

We will use this formula to show that we have $\nu(\chi) = 1$ for all irreducible characters χ of G. To do this, we first have to calculate the number of solutions x in G to the equation $x^2 = (0, 0)$.

Lemma 3. *The number of elements x in G that satisfy $x^2 = (0, 0)$ is*

$$q^5 + q^4 - q^3.$$

Proof. Suppose that $x = (u, a)$ in G satisfies $x^2 = (0, 0)$. Then since $(u, a)^2 = (0, f(u, u))$, we must have $f(u, u) = 0$. Therefore, we need to find how many vectors $u \in V$ satisfy $f(u, u) = 0$.

Since we are working over a finite field, it is easy to show that V has an orthonormal basis with respect to f. This means that there are basis vectors u_1 and u_2 of V, say, with

$$f(u_1, u_1) = f(u_2, u_2) = 1, \quad f(u_1, u_2) = 0.$$

Let u be any vector in V. We may write $u = \lambda u_1 + \mu u_2$ for unique elements λ and μ in $\mathbb{F}_{q^2}$. Then we have

$$f(u, u) = \lambda^{q+1} + \mu^{q+1}.$$

Thus $f(u, u) = 0$ if and only if $\lambda^{q+1} + \mu^{q+1} = 0$. We therefore need to count the number of ordered pairs (λ, μ) with $\lambda^{q+1} = \mu^{q+1}$, since we are working in a field of characteristic 2. Clearly, $\lambda = 0$ if and only if $\mu = 0$. On the other hand, if $\lambda \neq 0$, we have $(\lambda^{-1}\mu)^{q+1} = 1$. The number of elements z in $\mathbb{F}_{q^2}$ that satisfy $z^{q+1} = 1$ is $q + 1$. It follows that the number of ordered pairs of non-zero elements λ and μ satisfying $(\lambda^{-1}\mu)^{q+1} = 1$ is $(q^2 - 1)(q + 1)$. We deduce that the number of vectors $u \in V$ with $f(u, u) = 0$ is $(q^2 - 1)(q + 1) + 1 = q^3 + q^2 - q$.

Finally, the number of $x = (u, a)$ in G with $f(u, u) = 0$ is $q^2(q^3 + q^2 - q)$. □

Theorem 1. *We have $\nu(\chi) = 1$ for each irreducible complex character χ of G.*

Proof. It follows from Lemma 1 that

$$\sum_{\chi} \chi(1) = q^5 + (q^2 - q)q^2 = q^5 + q^4 - q^3,$$

where the sum extends over all irreducible complex characters χ of G. On the other hand, the Frobenius-Schur formula and Lemma 3 show that

$$\sum_{\chi} \nu(\chi)\chi(1) = q^5 + q^4 - q^3.$$

Since we have $\nu(\chi) \leq 1$, these equalities imply that $\nu(\chi) = 1$ for all irreducible complex characters χ. □

3. Isometries And Isotropic Subspaces

In this section, we consider isometries of V with respect to f and relate them to automorphisms of G.

We say that a non-zero $\mathbb{F}_{q^2}$-linear function $\sigma : V \to V$ is an *isometry* of V with respect to f if

$$f(\sigma(u), \sigma(v)) = f(u, v)$$

for all u and v in V. It is easy to see that, as f is non-degenerate, any isometry of V is invertible, and the inverse is also an isometry. The set of all isometries of V forms a group under composition, called the unitary group of degree 2 over $\mathbb{F}_{q^2}$ and denoted as $\mathrm{U}_2(\mathbb{F}_q)$.

Given an isometry σ of V with respect to f, we define $\hat{\sigma} : G \to G$ by setting

$$\hat{\sigma}(u, a) = (\sigma(u), a)$$

for all (u, a) in G. It is straightforward to see that $\hat{\sigma}$ is an automorphism of G which fixes $Z(G)$ elementwise. Furthermore, the mapping $\sigma \to \hat{\sigma}$ is a monomorphism. Thus, we see that $\mathrm{U}_2(\mathbb{F}_q)$ is embedded as a subgroup of the automorphism group of G.

We say that a non-zero vector u in V is *isotropic* if $f(u, u) = 0$. Suppose then that u is isotropic and let U be the one-dimensional $\mathbb{F}_{q^2}$-subspace spanned by u. It is easy to see that if v and w are any two (not necessarily different) elements in U, then $f(v, w) = 0$. We say that U is a *totally isotropic* subspace of V.

We note the following consequence of the proof of Lemma 3.

Lemma 4. *There are exactly $q + 1$ one-dimensional totally isotropic subspaces in V.*

Clearly, if σ is an isometry of V with respect to f and U is a one-dimensional totally isotropic subspace of V, $\sigma(U)$ is also totally isotropic. Thus $\mathrm{U}_2(\mathbb{F}_q)$ permutes the totally isotropic one-dimensional subspaces. We will show next that there is a cyclic subgroup of $\mathrm{U}_2(\mathbb{F}_q)$ of order $q + 1$ which transitively permutes these totally isotropic subspaces.

Let μ be a generator of the multiplicative group of non-zero elements of $\mathbb{F}_{q^2}$ and let $\lambda = \mu^{q-1}$. Then λ has order $q + 1$ in the multiplicative group. Using this notation, we have the transitivity property described above.

Lemma 5. *There is a cyclic subgroup of order $q + 1$ in $U_2(\mathbb{F}_q)$ which transitively permutes the one-dimensional totally isotropic subspaces.*

Proof. Let u_1 and u_2 be the orthonormal basis vectors of V previously introduced. Define $\sigma : V \to V$ by

$$\sigma(\alpha u_1 + \beta u_2) = \alpha\lambda u_1 + \beta u_2$$

for all α and β in $\mathbb{F}_{q^2}$. We claim that σ is an isometry of V. For we have

$$\begin{aligned} f(\sigma(u_1), \sigma(u_1)) &= \lambda\lambda^q f(u_1, u_1) = \lambda^{q+1} f(u_1, u_1) = f(u_1, u_1) \\ f(\sigma(u_1), \sigma(u_2)) &= f(\lambda u_1, u_2) = 0 = f(u_1, u_2) \\ f(\sigma(u_2), \sigma(u_2)) &= f(u_2, u_2) \end{aligned}$$

and these equations are enough to prove the desired claim.

Since $\sigma(u_1) = \lambda u_1$, $\sigma(u_2) = u_2$, and λ has order $q+1$, it is clear that σ has order $q+1$. We let H be the cyclic subgroup of $\mathrm{U}_2(\mathbb{F}_q)$ generated by σ. We note then that for any integer r satisfying $1 \leq r < q + 1$, σ has two different eigenvalues λ^r and

1, and precisely two different one-dimensional eigenspaces, namely, the subspaces U_1 and U_2, say, spanned by the eigenvectors u_1 and u_2. Moreover, U_1 and U_2 are not totally isotropic. Thus, no non-identity power of σ fixes a one-dimensional totally isotropic subspace of V and it follows that H acts transitively on these $q+1$ subspaces. □

Suppose next that U is one of the one-dimensional totally isotropic subspaces just considered. We set

$$\hat{U} = \{(u, 0) : u \in U\}.$$

Then $\hat{U}$ is a subset of G and since $f(u, v) = 0$ for all u and v in U, it is easy to see that $(u, 0)(v, 0) = (u + v, 0) = (v, 0)(u, 0)$. Thus $\hat{U}$ is an elementary abelian subgroup of G of order q^2. Corresponding to the $q+1$ totally isotropic subspaces of V, we obtain $q + 1$ elementary abelian subgroups of G of order q^2, which intersect trivially in pairs. Furthermore, if $\hat{\sigma}$ is the automorphism of G induced by the isometry σ, the cyclic subgroup of order $q+1$ generated by $\hat{\sigma}$ transitively permutes these $q + 1$ elementary abelian subgroups.

4. Action of Automorphisms on the Irreducible Representations of G

Let χ be an irreducible complex character of G of degree q^2. We have shown in Theorem 1 that χ is the character of a representation, X, say, of G defined on the real vector space $\mathbb{R}^{q^2}$. We may assume that X preserves a positive definite symmetric bilinear form, B, say, defined on $\mathbb{R}^{q^2} \times \mathbb{R}^{q^2}$.

Let τ be an isometry of V. We define the conjugate representation X^τ by

$$X^\tau(x) = X(\hat{\tau}(x))$$

for all $x \in G$. Since $\hat{\tau}$ fixes the elements of $Z(G)$ pointwise, and the character of X vanishes outside $Z(G)$, it is clear that X^τ also has character χ. It follows that X and X^τ are equivalent representations and thus there exists an automorphism $D(\tau)$, say, of $\mathbb{R}^{q^2}$ satisfying

$$X^\tau(x) = X(\hat{\tau}(x)) = D(\tau)X(x)D(\tau)^{-1}$$

for all $x \in G$.

Let us now take τ to be the isometry σ of order $q + 1$ described in Lemma 5 and let $D = D(\sigma)$. Then we have the following result.

Lemma 6. *Adopt the notation introduced above. Then multiplying D by a non-zero real scalar if necessary, we may arrange for D to satisfy $D^{q+1} = I$ and to preserve the bilinear form B.*

Proof. It is straightforward to verify that

$$D^i X(x) D^{-i} = X(\hat{\sigma}^i(x))$$

for each integer i and hence

$$D^{q+1} X(x) D^{-(q+1)} = X(x)$$

for all $x \in G$, since $\hat{\sigma}$ has order $q+1$. Therefore, D^{q+1} commutes with all $X(x)$ and hence is a real scalar multiple of the identity by Schur's Lemma, say $D^{q+1} = aI$, where a is a real number.

Taking determinants in the equality $D^{q+1} = aI$, we obtain

$$(\det D)^{q+1} = a^{q^2}.$$

We accordingly set $D_1 = (\det D)^{-1}a^{q-1}D$, so that D_1 is a scalar multiple of D, and calculate that

$$D_1^{q+1} = (\det D)^{-(q+1)}a^{(q^2-1)}D^{q+1} = a^{-1}D^{q+1} = I.$$

Thus, if we replace D by D_1, we obtain $D^{q+1} = I$.

Finally, let u and v be arbitrary elements of $\mathbb{R}^{q^2}$. We define a new symmetric bilinear form B' on $\mathbb{R}^{q^2} \times \mathbb{R}^{q^2}$ by setting

$$B'(u, v) = B(Du, Dv).$$

We intend to show that $B' = B$, implying that D preserves B. Now

$$\begin{aligned} B'(X(x)u, X(x)v) &= B(DX(x)u, DX(x)v) = B(X(\hat{\sigma}(x))Du, X(\hat{\sigma}(x))Dv) \\ &= B(Du, Dv) = B'(u, v) \end{aligned}$$

for all $x \in G$. This equality implies that X also preserves B'. But B is determined up to a scalar multiple of the identity, by Schur's Lemma, since X is irreducible. It follows that $B' = cB$ for some non-zero real number c. We note that c is positive, since B, and hence B', are positive definite. Thus we have

$$B(Du, Dv) = cB(u, v)$$

for all u and v in $\mathbb{R}^{q^2}$.

Taking powers, we obtain

$$B(D^{q+1}u, D^{q+1}v) = B(u, v) = c^{q+1}B(u, v)$$

and we deduce that $c^{q+1} = 1$. Since c is real and positive, it follows that $c = 1$. We deduce that D preserves B, as claimed.

□

5. Construction Of Real Mutually Unbiased Bases As Powers

Let U be a one-dimensional totally isotropic subspace of V and let $\hat{U}$ be the corresponding elementary abelian subgroup of G of order q^2. Since $\chi(x) = 0$ for any non-identity element x of $\hat{U}$ and $\chi(1) = q^2 = |\hat{U}|$, it follows that X defines the regular representation of $\hat{U}$ by restriction. Now the irreducible characters of $\hat{U}$ have degree 1, since the group is abelian, and they take only the values ± 1, since each element has order 1 or 2. The underlying real vector space $\mathbb{R}^{q^2}$ then splits as the direct sum of q^2 one-dimensional subspaces which are each invariant under the action of $X(\hat{U})$. Each of these one-dimensional subspaces determines a different irreducible (linear) character of $\hat{U}$.

As the elements of $X(\hat{U})$ preserve the underlying symmetric bilinear form B, the one-dimensional subspaces of $\mathbb{R}^{q^2}$ that we have described above are mutually orthogonal, and we may choose an orthonormal basis $Z_1, \ldots, Z_{q^2}$ of $\mathbb{R}^{q^2}$ such that

$$X(x)Z_i = \lambda_i(x)Z_i$$

for all $x \in \hat{U}$ and $1 \leq i \leq q^2$. Here, λ_i is an irreducible character of $\hat{U}$, with $\lambda_i(x) = \pm 1$, and $\lambda_1, \ldots, \lambda_{q^2}$ constitute all the irreducible characters of $\hat{U}$.

Now recall the isometry σ of V of order $q+1$ whose powers transitively permute the one-dimensional totally isotropic subspaces and let $\hat{\sigma}$ be the corresponding automorphism of G. We obtain $q+1$ subgroups $\hat{\sigma}^j(\hat{U})$, $0 \leq j \leq q$, that are all elementary abelian. It follows from the equation $D^j X(x) = X(\hat{\sigma}^j(x))D^j$ for all

$x \in \hat{U}$ that $D^j Z_1, \ldots, D^j Z_{q^2}$ is an orthonormal basis of eigenvectors of $X(\hat{\sigma}^j(\hat{U}))$ for $0 \leq j \leq q$. We thus obtain $q+1$ orthonormal bases of $\mathbb{R}^{q^2}$ from these subgroups.

The basic fact that we need to know is that these orthonormal bases are mutually unbiased. A proof of this fact is presented in the next theorem, in the context of properties of elementary abelian 2-subgroups of real orthogonal matrices. We follow the basic ideas laid out in the proof of Theorem 3.2 of [1].

Theorem 2. *Let $d = 2^k$, where k is a positive integer. Let M and N be elementary abelian 2-subgroups of $d \times d$ real orthogonal matrices, with $|M| = |N| = d$. Suppose that for all pairs $(x, y) \neq (I, I)$ in $M \times N$,*

$$\operatorname{tr}(xy) = 0,$$

where tr *denotes the trace of the matrix. Then any two orthonormal bases of $\mathbb{R}^d$ consisting of simultaneous eigenvectors of M and N, respectively, are mutually unbiased.*

Proof. We may assume that we have an orthonormal basis $X_1, \ldots, X_d$ of $\mathbb{R}^d$ such that

$$xX_i = \lambda_i(x)X_i$$

for all $x \in M$ and $1 \leq i \leq d$, where $\lambda_1, \ldots, \lambda_d$ are the irreducible characters of M. Likewise, we may assume that we have a second orthonormal basis $Y_1, \ldots, Y_d$ such that

$$yY_i = \mu_i(y)Y_i$$

for all $y \in N$ and $1 \leq i \leq d$, where $\mu_1, \ldots, \mu_d$ are the irreducible characters of N.

Consider now, for a given element x of M, the $d \times d$ symmetric matrix

$$X = \sum_{i=1}^{d} \lambda_i(x) X_i X_i^T.$$

Since the vectors $X_1, \ldots, X_d$ are orthonormal, we have

$$XX_j = \sum_{i=1}^{d} \lambda_i(x)(X_i X_i^T)X_j = \sum_{i=1}^{d} \lambda_i(x) X_i (X_i^T X_j) = \lambda_j(x) X_j$$

for any X_j. Thus X acts on the basis in an identical manner to x and we deduce that $X = x$. In the same way, we have

$$y = \sum_{i=1}^{d} \mu_i(y) Y_i Y_i^T$$

for any y in N.

Taking x and y as above, we obtain

$$xy = \sum_{1 \leq i,j \leq d} \lambda_i(x)\mu_j(y) X_i X_i^T Y_j Y_j^T,$$

and then we may take traces to obtain

$$\operatorname{tr}(xy) = \sum_{1 \leq i,j \leq d} \lambda_i(x)\mu_j(y) \operatorname{tr}(X_i X_i^T Y_j Y_j^T).$$

However, by simple properties of the trace function,

$$\operatorname{tr}(X_i X_i^T Y_j Y_j^T) = \operatorname{tr}(X_i^T Y_j Y_j^T X_i) = (X_i^T Y_j)^2$$

and thus we have

$$\mathrm{tr}(xy) = \sum_{1\le i,j\le d} \lambda_i(x)\mu_j(y)(X_i^T Y_j)^2.$$

Now by hypothesis, $\mathrm{tr}(xy) = 0$ and hence

$$0 = \sum_{1\le i,j\le d} \lambda_i(x)\mu_j(y)(X_i^T Y_j)^2,$$

unless $x = y = I$. When $x = y = I$, we have

$$d = \sum_{1\le i,j\le d} (X_i^T Y_j)^2.$$

As (x, y) runs over the elements of the direct product $M \times N$, we obtain a system of d^2 linear equations in the d^2 unknowns $(X_i^T Y_j)^2$, $1 \le i, j \le d$. The coefficient matrix of the system is the $d^2 \times d^2$ matrix C, say, which is the character table of $M \times N$. Taking (I, I) as the first element of $M \times N$, all entries of the first row of C equal 1, and C satisfies $CC^T = d^2 I$.

Our system of equations takes the form

$$\Psi = C\Phi,$$

where Φ is the column vector of size d^2, whose entries are the numbers $(X_i^T Y_j)^2$, and Ψ has its first entry equal to d, and all other entries equal to 0. Then we have

$$d^2\Phi = C^T\Psi,$$

and, since all entries of the first column of C^T are 1, we deduce that all entries of Φ equal $1/d$. It follows that

$$(X_i^T Y_j)^2 = \frac{1}{d}$$

for all i and j, and this equality implies that the bases are mutually unbiased.

□

Corollary 1. *Assume the notation previously introduced. Let $X_1, \ldots, X_{q^2}$ be an orthonormal basis of $\mathbb{R}^{q^2}$ consisting of simultaneous eigenvectors of the elements of $X(\hat{U})$. Then, with respect to this basis, the $q + 1$ powers D, D^2, $\ldots$, $D^{q+1} = I$, define $q + 1$ mutually unbiased bases of $\mathbb{R}^{q^2}$.*

We note that the fact that the powers of D define mutually unbiased bases means that the rescaled matrices qD, qD^2, $\ldots$, qD^q are q different real Hadamard matrices. The existence of these Hadamard matrices defined by the powers of a single orthogonal matrix gives a partial answer to the question raised in Section 5 of the author's earlier paper.

6. Diagonal Elements

Choosing the orthonormal basis of $\mathbb{R}^{q^2}$ that consists of simultaneous eigenvectors of the elements of $X(\hat{U})$, we may consider D and its powers to be explicit real orthogonal matrices, and we shall adopt this point of view. The entries of the non-identity powers of the matrix D are then all $\pm 1/q$. We intend to show that in fact the diagonal elements are all $-1/q$. If we could describe D explicitly, this would be a trivial observation, but we suspect that it is difficult to describe D, and consequently we will proceed indirectly by employing some considerations of a character-theoretic nature.

Let S be the cyclic subgroup of the automorphism group of G generated by $\hat{\sigma}$. D determines a representation of S of degree q^2, in which an element $\hat{\sigma}^i$ is mapped to D^i. Let θ denote the character of this representation. Thus

$$\theta(\hat{\sigma}^i) = \mathrm{tr}(D^i).$$

Since the trace of a matrix is the sum of its diagonal entries, we have the following estimate for the values taken by θ.

Lemma 7. *For $1 \leq i \leq q$, we have*

$$-q \leq \theta(\hat{\sigma}^i) \leq q.$$

Proof. The lower estimate occurs if all diagonal elements of D^i are $-1/q$ and the upper estimate occurs if all diagonal elements are $1/q$. □

We shall speak of the subspace of fixed elements of all the powers of D acting on $\mathbb{R}^{q^2}$ as the fixed point space of S in its representation. As is well known, the dimension of this fixed point space is the inner product of the principal or trivial character of S with θ, and thus equals

$$\frac{1}{q+1}\left(q^2 + \sum_{i=1}^{q} \mathrm{tr}(D^i)\right).$$

The next lemma relates the triviality of the fixed point space of S with the stated diagonal property of the powers of D.

Lemma 8. *The fixed point space of S is trivial if and only if all diagonal elements of each non-identity element D^i, $1 \leq i \leq q$, equal $-1/q$.*

Proof. In terms of the character θ, the fixed point space of S is trivial if and only if

$$q^2 + \sum_{i=1}^{q} \theta(\hat{\sigma}^i) = 0.$$

However, Lemma 7 implies that

$$\sum_{i=1}^{q} \theta(\hat{\sigma}^i) \geq -q^2,$$

with equality occurring if and only if each diagonal element of D^i equals $-1/q$, for $1 \leq i \leq q$. The proof of the lemma is now clear.

□

One approach to showing that the fixed point subspace of S is trivial is clearly to calculate the character values of θ and we believe this is possible by applying some character theory. We will take a more oblique approach by examining properties of the fixed point space.

We begin by obtaining an upper bound for the dimension of the fixed point space.

Lemma 9. *The dimension of the fixed point space of S is at most $2q - 2$.*

Proof. Let t be the dimension of the fixed point subspace. As we have seen,

$$t = \frac{1}{q+1}\left(q^2 + \sum_{i=1}^{q} \mathrm{tr}(D^i)\right).$$

Now Lemma 7 implies that

$$\sum_{i=1}^{q} \mathrm{tr}(D^i) \leq q^2$$

and the upper bound for t follows from this inequality. □

We return to a study of the isometry σ of V that gives rise to $\hat{\sigma}$. We defined σ in such a way that it fixes the vector u_2 that is part of the orthonormal basis of V. For simplicity, we shall denote this vector in this section by u. Let C denote the subgroup of G consisting of all elements of the form

$$(\lambda u, \mu),$$

where λ and μ range independently over $\mathbb{F}_{q^2}$. It is clear that C has order q^4.

The following lemma is easy to prove by elementary considerations.

Lemma 10. *Let λ be a non-zero element of $\mathbb{F}_{q^2}$. Then the centralizer of $(\lambda u, \mu)$ in C has order q^3. The center $Z(C)$ of C coincides with $Z(G)$. The commutator subgroup of C consists of all elements $(0, \kappa)$, where $\kappa \in \mathbb{F}_q$, and hence has order q. The involutions in C are the elements of the form $(0, \mu)$, where $\mu \neq 0$, and there are exactly $q^2 - 1$ of them.*

The next result relating to the characters of C is a close analogue of earlier results obtained for G.

Lemma 11. *The group C has exactly q^3 irreducible complex characters of degree 1. All other irreducible complex characters of C have degree q and there are $q^2 - q$ of them. The irreducible characters of degree q are real-valued, vanish outside $Z(C)$, and each has Frobenius-Schur indicator equal to -1.*

Proof. Since the commutator subgroup C' has order q, C has exactly $|C : C'| = q^3$ different irreducible characters of degree 1. The argument used to prove Lemma 1 implies that, as each non-central element x in C has a centralizer in C of order q^3, $\psi(x) = 0$ for any irreducible character ψ of C of degree greater than 1. Then, again arguing as in the proof of Lemma 1, we deduce that $\psi(1) = q$, and that there are $q^2 - q$ different irreducible characters of degree q.

Concerning the reality of the characters of C, we argue as follows. C' consists of all elements of the form $(0, \kappa)$, where $\kappa \in \mathbb{F}_q$. Now given any element $x = (\lambda u, \mu)$, we have $x^2 = (0, \lambda\lambda^q) \in C'$. This implies that C/C' is an elementary abelian 2-group and thus each character of C of degree 1 is real valued and trivially has Frobenius-Schur indicator equal to 1. Since C contains exactly $q^2 - 1$ involutions, the Frobenius-Schur involution formula implies that

$$\sum_{\psi} \nu(\psi)\psi(1) = q^2 - q^3,$$

where the sum extends over all irreducible characters ψ of C of degree greater than one. Since there are $q^2 - q$ such characters, and they all have degree q, we deduce that $\nu(\psi) = -1$ for each such character ψ, as required.

□

For our next lemma, we introduce some standard notation. Given any character θ, say, of G, we let θ_C (or $(\theta)_C$), denote its restriction to C.

Lemma 12. *Given any irreducible character ψ of C of degree q, there is a unique irreducible character χ_ψ of G of degree q^2 that satisfies $(\chi_\psi)_C = q\psi$. There is a one-to-one correspondence between the irreducible characters of G of degree q^2 and those of C of degree q, determined by restriction of characters to C*

Proof. Let ψ be an irreducible character of C of degree q. Since ψ vanishes outside $Z(C)$, and $Z(C) = Z(G)$, it follows from the definition of induced characters that

$$\psi^G(x) = 0$$

if $x \notin Z(G)$, and

$$\psi^G(z) = q^2\psi(z)$$

if $z \in Z(G)$. Thus

$$(\psi^G)_C = q^2\psi.$$

We deduce that the restriction to C of any irreducible constituent χ, say, of ψ^G is a multiple of ψ. Thus, since necessarily $\chi(1) = q^2$, we see that $\chi_C = q\psi$. The Frobenius reciprocity theorem implies that we have the equality

$$(\psi^G, \chi) = (\psi, \chi_C) = q$$

of inner products. This implies that χ is a constituent of ψ^G with multiplicity q. However, since $\psi^G(1) = q^3 = q\chi(1)$, a comparison of degrees shows that we must have $\psi^G = q\chi$.

We can now write $\chi = \chi_\psi$ and associate χ_ψ with ψ. Since there are equal numbers (q^2-q) of irreducible characters of G of degree q^2 and irreducible characters of C of degree q, we deduce that we have established a one-to-one correspondence between the two sets of characters. □

Corollary 2. *The fixed point subspace of S acting on $\mathbb{R}^{q^2}$ is trivial.*

Proof. Let U denote the fixed point subspace of S acting on $\mathbb{R}^{q^2}$. Since S fixes all the elements of the subgroup C of G, it follows that D and its powers commute with the elements $X(c)$, as c runs over C. It is straightforward to see from this that U is invariant under the action of $X(C)$. U is thus expressible as the direct sum of various irreducible C-submodules.

Now Lemma 12 shows that $\mathbb{R}^{q^2}$ is a direct sum of q isomorphic irreducible C-submodules, each of dimension q. Thus U is the direct sum of irreducible submodules of C, each of dimension q. We also have the dimension bound $\dim U \leq 2q-2$ from Lemma 9. We deduce from this upper bound that, if U is non-trivial, it must be an irreducible C-submodule of $\mathbb{R}^{q^2}$ of dimension q. This however is impossible, since U defines then an irreducible real representation of C of degree q, whereas we have shown in Lemma 11 that no irreducible representation of degree q of C can be defined over the real numbers. It follows that U is trivial, as required. □

Corollary 3. *The diagonal elements of each matrix D^i are $-1/q$, for $1 \leq i \leq q$.*

References

[1] S. Bandyopadhyay, P. O. Boykin, V. Roychowdhury and F. Vatan, *A new proof for the existence of mutually unbiased bases*, Algorithmica **34** (2002), 512-528.

[2] P. O. Boykin, M. Sitharam, M. Tarifi and P. Wocjan, *On the maximal number of real mutually unbiased bases*, arXiv.org e-Print archive quant-ph/0502024

[3] P. J. Cameron and J. J. Seidel, *Quadratic forms over $GF(2)$*, Indag. Math **25** (1973), 1-8.

[4] R. Gow, *Generation of mutually unbiased bases as powers of a unitary matrix in 2-power dimensions*, arXiv.org e-Print archive math/0703333.

[5] I. M. Isaacs, *Character Theory of Finite Groups*, Academic Press, New York, 1976.